



## Post-Traumatic Stress Disorder

### *April 2006*

1: Schwartz AC, Bradley R, Penza KM, Sexton M, Jay D, Haggard PJ, Garlow SJ, Ressler KJ.

Pain medication use among patients with posttraumatic stress disorder.  
Psychosomatics. 2006 Mar-Apr; 47(2):136-42.

PMID: 16508025 [PubMed - indexed for MEDLINE]

2: Hoge CW, Auchterlonie JL, Milliken CS.

Mental health problems, use of mental health services, and attrition from military service after returning from deployment to Iraq or Afghanistan.  
JAMA. 2006 Mar 1; 295(9):1023-32.

PMID: 16507803 [PubMed - indexed for MEDLINE]

3: Budur K, Falcone T, Franco K.

Diagnosing and managing posttraumatic stress disorder.  
Cleve Clin J Med. 2006 Feb; 73(2):121-2, 125-6, 128-9. Review.  
PMID: 16478037 [PubMed - indexed for MEDLINE]

4: Pfefferbaum B, Tucker P, North CS, Jeon-Slaughter H, Kent AT, Schorr JK, Wilson TG, Bunch K.

Persistent physiological reactivity in a pilot study of partners of firefighters after a terrorist attack.

J Nerv Ment Dis. 2006 Feb; 194(2):128-31.

PMID: 16477192 [PubMed - indexed for MEDLINE]

5: Briere J.

Dissociative symptoms and trauma exposure: specificity, affect dysregulation, and posttraumatic stress.

J Nerv Ment Dis. 2006 Feb; 194(2):78-82.

PMID: 16477184 [PubMed - indexed for MEDLINE]

6: Pitman RK.

Combat effects on mental health: the more things change, the more they remain the same.

Arch Gen Psychiatry. 2006 Feb; 63(2):127-8. No abstract available.

PMID: 16461853 [PubMed - indexed for MEDLINE]